



PERFIL DEL PACIENTE

INFORMACIÓN DEL PACIENTE

FECHA: _____

NOMBRE: _____

DIRECCIÓN: _____

CIUDAD: _____ CODIGO POSTAL: _____

TELÉFONO/ CASA: _____ TELÉFONO/ MÓVIL: _____

TELÉFONO/TRABAJO: _____

GENDER: HOMBRE MUJER OTRO

FECHA DE NACIMIENTO: _____

¿Cómo se entero de nosotros? _____



REGISTRO DE VACUNACIÓN DE CALIFORNIA – REGIÓN IV

DECLARACIÓN DE REVELACIÓN: PARA PACIENTES O PADRES DE FAMILIA

Somos miembros autorizados del Registro de Vacunación de California – Región IV. El Registro de Vacunación permite que se compartan datos electrónicos de vacunación y de pruebas de detección de tuberculosis de usted y de su hijo independientemente de donde se hayan administrado. Usted y su doctor pueden ver en cualquier momento qué vacunas les dieron y qué pruebas de tuberculosis les hicieron a usted o a su hijo y qué vacunas se tienen que dar ahora. Ayudará a usted y a su doctor a protegerlos a usted y a su hijo contra enfermedades serias como la poliomielitis, tos ferina, sarampión y meningitis.

La información del Registro de Vacunación es confidencial. Su información **SOLO** se compartirá con (a.) profesionales de la salud (o sea doctores, clínicas u hospitales), para ayudar a decidir qué vacunas necesitan usted o su hijo; para llamarlo por teléfono o enviarle un recordatorio de que le toca recibir una vacuna; y, contar cuántos pacientes están o no al día con sus vacunas, (b.) escuelas o guarderías de niños, para ayudar a demostrar que usted o su hijo han recibido las vacunas requeridas para ingresar, (c.) las clínicas WIC, para informarles si a su hijo le hace falta una dosis de una vacuna, (d.) planes de atención de la salud, para ayudar a procesar pagos de seguros (e.) el Intercambio de Información de Salud del Condado de San Joaquin (SJC HIE, en inglés) y (f.) la División de Vacunación del Departamento de Salud Pública del Estado de California.

La siguiente es la información que el Registro de Vacunación mantendrá sobre usted o su hijo:

- Nombre y fecha de nacimiento
- Nombres de los padres de familia o tutores
- Sexo (masculino o femenino)
- Dirección y teléfono actuales (solo los profesionales de la salud pueden ver esta información)
- Tipos de vacunas y pruebas de tuberculosis administradas y las fechas
- Reacciones serias a vacunas o pruebas de tuberculosis, si corresponde
- Otra información limitada que pueda ayudar a identificar correctamente a usted o a su hijo

Derechos de pacientes y padres

Tiene derecho legal a pedir lo siguiente:

- Que los datos en el registro de vacunaciones y pruebas de tuberculosis de usted y de su hijo se compartan solo con su doctor*
- Que el Registro no le envíe recordatorios de citas
- Ver una copia de los datos de vacunación y de pruebas de tuberculosis de usted y de su hijo que figuren en el Registro
- Que le digan quiénes han visto los datos en el Registro o que su doctor corrija cualquier error

Si DESEA que sus datos o los de su hijo estén en el Registro, no haga nada. Eso es todo.

Si NO DESEA que el consultorio de su doctor comparta sus datos de vacunación e información sobre sus pruebas de tuberculosis con otros usuarios del Registro, pida en el consultorio de su doctor que le den un “Formulario de rechazo”.

Para más información sobre sus derechos, póngase en contacto con su profesional de la salud.

Para más información sobre el Registro de vacunación, llame al (209) 468-2292.

**Por ley, si hay una emergencia de salud pública los funcionarios de salud pública pueden ver el Registro.*

Cuestionario de contraindicaciones para vacunación de niños y adolescentes

NOMBRE DEL PACIENTE _____

FECHA DE NACIMIENTO ____/____/____
mes día año

A los padres/tutores: Las siguientes preguntas nos ayudarán a determinar cuáles vacunas le podremos administrar a su hijo hoy. Si responde “sí” a alguna pregunta, no necesariamente significa que no se debe vacunar a su hijo. Simplemente quiere decir que hay que hacerle más preguntas. Si alguna pregunta no está clara, solicítele a su proveedor de atención médica que se la explique.

	sí	no	no sé
1. ¿El niño está enfermo hoy?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. ¿El niño es alérgico a algún medicamento, alimento, componente de vacunas o al látex?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3. ¿El niño ha tenido alguna reacción seria a una vacuna en el pasado?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4. ¿El niño tiene algún problema de salud crónico en los pulmones, el corazón o los riñones, o sufre de enfermedad metabólica (p. ej., diabetes), asma, un trastorno de la sangre, no tiene bazo, tiene deficiencia de componentes del complemento, un implante coclear o derrame de líquido cefalorraquídeo? ¿Está en terapia de aspirina a largo plazo?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. Si el niño que va a ser vacunado tiene entre 2 y 4 años, ¿le ha dicho algún proveedor de atención médica que el niño tuvo sibilancia o asma en los últimos 12 meses?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6. Si el niño es un bebé, ¿le han dicho alguna vez que tuvo intususcepción?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
7. ¿El niño, uno de sus hermanos o uno de sus padres ha tenido convulsiones; o el niño ha tenido problemas cerebrales o algún otro problema del sistema nervioso?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
8. ¿El niño o un miembro de su familia tiene cáncer, leucemia, VIH/SIDA o cualquier otro problema del sistema inmunitario?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
9. ¿Uno de los padres, hermanos o hermanas del niño tiene algún problema en su sistema inmunitario?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
10. ¿En los últimos 3 meses, el niño ha tomado medicamentos que afectan el sistema inmunitario, tales como prednisona, otros esteroides o medicamentos contra el cáncer, o medicamentos para el tratamiento de la artritis reumatoide, la enfermedad de Crohn o la psoriasis, o tuvo tratamientos de radiación?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
11. Durante el año pasado, ¿el niño recibió una transfusión de sangre o de productos sanguíneos, o se le administró inmunoglobulina, gammaglobulina o algún medicamento antiviral?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
12. ¿La niña/adolescente está embarazada o hay alguna posibilidad de que quede embarazada durante el próximo mes?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
13. ¿Se le aplicó alguna vacuna al niño en las últimas 4 semanas?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

FORMA LLENADA POR _____ FECHA _____

FORMA REVISADA POR _____ FECHA _____

¿Trajo su cartilla de vacunación consigo? sí no

Es importante que tenga un registro personal de las vacunas de su hijo. Si no lo tiene, pídale al proveedor de atención médica de su hijo que le dé uno con todas las vacunas de su hijo. Guárdelo en un lugar seguro y llévelo con usted todas las veces que busque atención médica para su hijo. Su hijo necesitará este documento para ingresar a la guardería o a la escuela, para obtener empleo o para viajar al extranjero.

“Screening Checklist for Contraindications to Vaccines for Children and Teens”

Saint Paul, Minnesota • 651-647-9009 • www.immunize.org • www.vaccineinformation.org

www.immunize.org/catg.d/p4060-01.pdf • Item #P4060-01 Spanish (10/20)

Information for Healthcare Professionals about the Screening Checklist for Contraindications to Vaccines (Children and Teens)

Are you interested in knowing why we included a certain question on the screening checklist? If so, read the information below. If you want to find out even more, consult the references in **Notes** below.

NOTE: For supporting documentation on the answers given below, go to the specific ACIP vaccine recommendation found at the following website: www.cdc.gov/vaccines/hcp/acip-recs/index.html

1. Is the child sick today? [all vaccines]

There is no evidence that acute illness reduces vaccine efficacy or increases vaccine adverse events. However, as a precaution with moderate or severe acute illness, all vaccines should be delayed until the illness has improved. Mild illnesses (such as otitis media, upper respiratory infections, and diarrhea) are NOT contraindications to vaccination. Do not withhold vaccination if a person is taking antibiotics.

2. Does the child have allergies to medications, food, a vaccine component, or latex? [all vaccines]

An anaphylactic reaction to latex is a contraindication to vaccines that contain latex as a component or as part of the packaging (e.g., vial stoppers, prefilled syringe plungers, prefilled syringe caps). If a person has anaphylaxis after eating gelatin, do not administer vaccines containing gelatin. A local reaction to a prior vaccine dose or vaccine component, including latex, is not a contraindication to a subsequent dose or vaccine containing that component. For information on vaccines supplied in vials or syringes containing latex, see www.cdc.gov/vaccines-pubs/pinkbook/downloads/appendices/B/latex-table.pdf; for an extensive list of vaccine components, see www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/B/excipient-table-2.pdf. People with egg allergy of any severity can receive any recommended influenza vaccine (i.e., any IIV, RIV, or LAIV) that is otherwise appropriate for the patient's age and health status. With the exception of cIIIV and RIV (which do not contain egg antigen), people with a history of severe allergic reaction to egg involving any symptom other than hives (e.g., angioedema, respiratory distress), or who required epinephrine or another emergency medical intervention, the vaccine should be administered in a medical setting, such as a clinic, health department, or physician office; vaccine administration should be supervised by a healthcare provider who is able to recognize and manage severe allergic conditions.

3. Has the child had a serious reaction to a vaccine in the past? [all vaccines]

History of anaphylactic reaction (see question 2) to a previous dose of vaccine or vaccine component is a contraindication for subsequent doses. History of encephalopathy within 7 days following DTP/DTPa is a contraindication for further doses of pertussis-containing vaccine. There are other adverse events that might have occurred following vaccination that constitute contraindications or precautions to future doses. Under normal circumstances, vaccines are deferred when a precaution is present. However, situations may arise when the benefit outweighs the risk (e.g., during a community pertussis outbreak).

4. Does the child have a long-term health problem with lung, heart, kidney, or metabolic disease (e.g., diabetes), asthma, a blood disorder, no spleen, complement component deficiency, a cochlear implant, or a spinal fluid leak? Is he/she on long-term aspirin therapy? [MMR, MMRV, LAIV, VAR]

A history of thrombocytopenia or thrombocytopenic purpura is a precaution to MMR and MMRV vaccines. The safety of LAIV in children and teens with lung, heart, kidney, or metabolic disease (e.g., diabetes), or a blood disorder has not been established. These conditions, including asthma in children ages 5 years and older, should be considered precautions for the use of LAIV. Children with functional or anatomic asplenia, complement deficiency, cochlear implant, or CSF leak should not receive LAIV. Children on long-term aspirin therapy should not be given LAIV; instead, they should be given IIV. Children with CSF leak, anatomic or functional asplenia, or cochlear implant, or on long-term aspirin therapy should not be given LAIV; instead, they should be given IIV. Aspirin use is a precaution to VAR.

5. If the child to be vaccinated is 2 through 4 years of age, has a healthcare provider told you that the child had wheezing or asthma in the past 12 months? [LAIV]

Children ages 2 through 4 years who have had a wheezing episode within the past 12 months should not be given LAIV. Instead, these children should be given IIV.

6. If your child is a baby, have you ever been told that he or she has had intussusception? [Rotavirus]

Infants who have a history of intussusception (i.e., the telescoping of one portion of the intestine into another) should not be given rotavirus vaccine.

7. Has the child, a sibling, or a parent had a seizure; has the child had brain or other nervous system problem? [DTaP, Td, Tdap, IIV, LAIV, MMRV]

DTaP and Tdap are contraindicated in children who have a history of encephalopathy within 7 days following DTP/DTPa. An unstable progressive neurologic problem is a precaution to the use of DTaP and Tdap. For children with stable neurologic disorders (including seizures) unrelated to vaccination, or for children with a family history of seizures, vaccinate as usual (exception: children with a personal or family [i.e., parent or sibling] history of seizures generally should not be vaccinated with MMRV; they should receive separate MMR and VAR vaccines). A history of Guillain-Barré syndrome (GBS) is a consideration with the following: 1) Td/Tdap: if GBS has occurred within 6 weeks of a tetanus-containing vaccine and decision is made to continue vaccination, give Tdap instead of Td if no history of prior Tdap;

NOTE: For summary information on contraindications and precautions to vaccines, go to the ACIP's General Best Practice Guidelines for Immunization at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html

2) Influenza vaccine (IIV, LAIV, or RIV): if GBS has occurred within 6 weeks of a prior influenza vaccination, vaccinate with IIV if at high risk for severe influenza complications.

8. Does the child have cancer, leukemia, HIV/AIDS, or any other immune system problem? [LAIV, MMR, MMRV, RV, VAR]

Live virus vaccines (e.g., MMR, MMRV, VAR, RV, LAIV) are usually contraindicated in immunocompromised children. However, there are exceptions. For example, MMR is recommended for asymptomatic HIV-infected children who do not have evidence of severe immunosuppression. Likewise, VAR should be considered for HIV-infected children age 12 months through 8 years with age-specific CD4+ T-lymphocyte percentage at 15% or greater, or for children age 9 years or older with CD4+ T-lymphocyte counts of greater than or equal to 200 cell/ μ L. VAR should be administered (if indicated) to persons with isolated humoral immunodeficiency. Immunosuppressed children should not receive LAIV. Infants who have been diagnosed with severe combined immunodeficiency (SCID) should not be given a live virus vaccine, including RV. Other forms of immunosuppression are a precaution, not a contraindication, to RV. For details, consult ACIP recommendations (see references in **Notes** above).

9. Does the child have a parent, brother, or sister with an immune system problem? [MMR, MMRV, VAR]

MMR, VAR, and MMRV vaccines should not be given to a child or teen with a family history of congenital or hereditary immunodeficiency in first-degree relatives (i.e., parents, siblings) unless the immune competence of the potential vaccine recipient has been clinically substantiated or verified by a laboratory.

10. In the past 3 months, has the child taken medications that affect the immune system such as prednisone, other steroids, or anticancer drugs; drugs for the treatment of rheumatoid arthritis, Crohn's disease, or psoriasis; or had radiation treatments? [LAIV, MMR, MMRV, VAR]

Live virus vaccines (e.g., LAIV, MMR, MMRV, VAR) should be postponed until after chemotherapy or long-term high-dose steroid therapy has ended. For details and length of time to postpone, consult the ACIP statement. Some immune mediator and immune modulator drugs (especially the antitumor-necrosis factor agents adalimumab, infliximab, and etanercept) may be immunosuppressive. A comprehensive list of immunosuppressive immune modulators is available in CDC Health Information for International Travel (the "Yellow Book") available at wwwnc.cdc.gov/travel/yellowbook/2020/travelers-with-additional-considerations/immunocompromised-travelers. The use of live vaccines should be avoided in persons taking these drugs. To find specific vaccination schedules for stem cell transplant (bone marrow transplant) patients, see General Best Practice Guidelines for Immunization (referenced in **Notes** above). LAIV, when recommended, can be given only to healthy non-pregnant people ages 2 through 49 years.

11. In the past year, has the child received a transfusion of blood/blood products, or been given immune (gamma) globulin or an antiviral drug? [MMR, MMRV, LAIV, VAR]

Certain live virus vaccines (e.g., MMR, MMRV, LAIV, VAR) may need to be deferred, depending on several variables. Consult the most current ACIP recommendations (referenced in **Notes** above) for the most current information on intervals between antiviral drugs, immune globulin or blood product administration and live virus vaccines.

12. Is the child/teen pregnant or is there a chance she could become pregnant during the next month? [HPV, IPV, LAIV, MenB, MMR, MMRV, VAR]

Live virus vaccines (e.g., MMR, MMRV, VAR, LAIV) are contraindicated one month before and during pregnancy because of the theoretical risk of virus transmission to the fetus. Sexually active young women who receive a live virus vaccine should be instructed to practice careful contraception for one month following receipt of the vaccine. On theoretical grounds, IPV and MenB should not be given during pregnancy; however, it may be given if there is a risk of exposure. IIV and Tdap are both recommended during pregnancy. HPV vaccine is not recommended during pregnancy.

13. Has the child received vaccinations in the past 4 weeks? [LAIV, MMR, MMRV, VAR, yellow fever]

Children who were given either LAIV or an injectable live virus vaccine (e.g., MMR, MMRV, VAR, yellow fever) should wait 28 days before receiving another vaccination of this type (30 days for yellow fever vaccine). Inactivated vaccines may be given at the same time or at any spacing interval.

VACCINE ABBREVIATIONS

IIV = Live attenuated influenza vaccine	MMRV = MMR+VAR vaccine
HPV = Human papillomavirus vaccine	RIV = Recombinant influenza vaccine
IIV = Inactivated influenza vaccine	RV = Rotavirus vaccine
cIIIV = cell culture inactivated influenza vaccine	Td/Tdap = Tetanus, diphtheria, (acellular pertussis) vaccine
IPV = Inactivated poliovirus vaccine	VAR = Varicella vaccine
MMR = Measles, mumps, and rubella vaccine	



Registro para la Evaluación de Elegibilidad de Pacientes para el Programa de Vacunas para Niños (VFC)

Nombre del Paciente:	<i>Apellido</i>	<i>Primer</i>	Fecha de Hoy:
Fecha de Nacimiento: / /	Padre/Tutor	<i>Apellido</i>	<i>Primer</i>
Edad de Paciente:			
Idioma Principal del Paciente:	<input type="checkbox"/> Español <input type="checkbox"/> Inglés <input type="checkbox"/> Otra		
Género (por favor marque)	Hombre Mujer Otro		
Etnicidad (por favor marque)	blanco afroamericano/negro indio americano asiático Nativo de Alaska Nativo de Hawái/isleño del Pacífico		
Dirección:	Calle: Ciudad y Estado: Codigo postal:		
Número de teléfono:			
Correo electrónico:			
¿Tiene Medi-Cal?	<input type="checkbox"/> Sí	<input type="checkbox"/> No	
¿Tienes un seguro privado?	<input type="checkbox"/> Sí	<input type="checkbox"/> No	

- Este formulario documenta si el paciente mencionado anteriormente reúne los requisitos necesarios.
- El profesional de la salud debe mantener este expediente para cada niño que reúne los requisitos para VFC por un mínimo de tres (3) años y ponerlos a la disposición para la inspección a solicitud de las autoridades estatales o federales.
- El paciente, su padre/madre o profesional de la salud puede llenar este formulario (si él/ella es un menor emancipado [independiente] o tiene 18 años de edad).
- La evaluación de elegibilidad para el Programa VFC y la documentación del estado de elegibilidad debe hacerse durante cada visita de vacunación para asegurar que el estado de elegibilidad no ha cambiado.
- No se requiere la verificación de respuestas.



ESTA SECCIÓN DEBE SER COMPLETADA POR PERSONAL DE LA OFICINA

La Verificación del Estado de Elegibilidad

Fecha de la evaluación	VFC Elegible				Not VFC Eligible
	Elegible para Medi-Cal o CHPD	No tiene seguro médico	Nativo Americano/ Nativo de Alaska	El seguro médico no cubre el costo de la(s) vacuna(s)	Asegurado (El paciente tiene Seguro)

El paciente califica para la inmunización a través del Programa VFC porque él/ella o sus padres/tutores declaran que el niño tiene 18 años de edad o menos Y:

Elija solo uno de los siguientes.

(Nota: si un niño cumple con dos o más de los requisitos de elegibilidad, elija el primero que corresponda).

- Es elegible para Medi-Cal o Salud Infantil y Prevención de Discapacidades (CHDP); o
- Esta sin seguro (no tiene seguro médico privado); o
- Es Indio Americano o Nativo de Alaska o
- El seguro de salud no cubre vacunas (solo en centros de salud rurales y calificados federalmente).

El paciente no califica para la inmunización a través del Programa VFC porque tiene seguro médico que paga las vacunas.

Comentarios: